

## Urecholine



**Generic Urecholine** is a diuretic, which stimulates your bladder. It is used to cure urinary retention, in other words difficulty urinating, which may be experienced after surgery, after delivery etc.

Active ingredient: [Bethanechol](#)

**Brand(s):**Urivoid / Macpee  
**Manufacturer:** Cipla Limited / Intas Pharmaceuticals Ltd  
**Disease(s):**Urinary Retention / Difficulty Urinating

### Urecholine 25mg

|            | Per pill | Per Pack                         | Order                       |
|------------|----------|----------------------------------|-----------------------------|
| 10 tablet  | \$ 3.00  | \$ 29.95                         | <a href="#">Add to cart</a> |
| 20 tablet  | \$ 2.50  | \$ 49.95<br><del>\$59.90</del>   | <a href="#">Add to cart</a> |
| 30 tablet  | \$ 2.00  | \$ 59.95<br><del>\$89.85</del>   | <a href="#">Add to cart</a> |
| 60 tablet  | \$ 1.75  | \$ 104.95<br><del>\$179.70</del> | <a href="#">Add to cart</a> |
| 90 tablet  | \$ 1.50  | \$ 134.95<br><del>\$269.55</del> | <a href="#">Add to cart</a> |
| 120 tablet | \$ 1.25  | \$ 149.95<br><del>\$359.40</del> | <a href="#">Add to cart</a> |

### Bethanechol tablet

#### What is this medicine?

BETHANECHOL stimulates your bladder to empty.

Bethanechol is used to treat urinary retention (difficulty urinating), which may occur after surgery, after delivering a baby, and in other situations.

Bethanechol may also be used for purposes other than those listed in this medication guide.

#### What is the most important information I should know about Urecholine (bethanechol)?

Take bethanechol on an empty stomach 1 hour before or 2 hours after meals to prevent nausea and vomiting.

Bethanechol may cause dizziness or fainting, especially when you rise from a sitting or lying position. Rise slowly to avoid becoming dizzy, falling, or hurting yourself.

Use caution when driving, operating machinery, or performing other hazardous activities. Bethanechol may cause dizziness. If you experience dizziness, avoid these activities.

#### What should I discuss with my healthcare provider before taking Urecholine (bethanechol)?

Before taking this medication, tell your doctor if you

- have uncontrolled hyperthyroidism (an overactive thyroid);
- have stomach ulcers;

