

Phenergan Syrup



Generic Phenergan belongs to a group of drugs called phenothiazines. It works by changing the actions of chemicals in your brain. Promethazine (the active ingredient) also acts as an antihistamine. It blocks the effects of the naturally occurring chemical histamine in your body. Phenergan is used to treat allergy symptoms such as itching, runny nose, sneezing, itchy or watery eyes, hives, and itchy skin rashes. Phenergan also prevents motion sickness, and treats nausea and vomiting or pain after surgery. It is also used as a sedative or sleep aid.

Active ingredient: [Promethazine](#)

Brand(s):Phenergan Syrup

Manufacturer: Abbott Laboratories

Disease(s):Allergy / Motion Sickness / Nausea

Phenergan Syrup 100ml

	Per bottle	Per Pack	Order
1 bottles	\$ 9.95	\$ 9.95	Add to cart
2 bottles	\$ 7.47	\$ 14.95 \$19.90	Add to cart
3 bottles	\$ 6.65	\$ 19.95 \$29.85	Add to cart
4 bottles	\$ 5.74	\$ 22.95 \$39.80	Add to cart
5 bottles	\$ 4.99	\$ 24.95 \$49.75	Add to cart

Promethazine syrup

What is this medicine?

PROMETHAZINE is an antihistamine. It relieves moderate to severe allergic reactions, reduces or prevents nausea and vomiting from illness or motion sickness, helps to make you sleep before surgery, and may help with pain relief or nausea after surgery.

What should I tell my health care provider before I take this medicine?

They need to know if you have any of these conditions:

- asthma or congestive lung disease
- diabetes
- glaucoma
- high blood pressure or heart disease
- kidney disease
- liver disease
- prostate trouble
- pain or difficulty passing urine
- seizures (convulsions)
- an unusual or allergic reaction to promethazine or phenothiazines, sulfites, other medicines, foods, dyes, or preservatives
- pregnant or trying to get pregnant
- breast-feeding

How should I use this medicine?

Use Phenergan as directed by your doctor. Take Promethazine by mouth with or without food. If stomach upset occurs, take with food to reduce stomach irritation. If you are using it for motion sickness, take a dose at least 30 to 60 minutes before you begin traveling.

