Lopid



Generic Lopid is used for treating high blood cholesterol and trialyceride.

Active ingredient: Gemfibrozil

Brand(s):Lopid
Manufacturer: Pfizer Inc.
Disease(s):Hypercholesterolemia

Lopid 300mg

| | Per pill | Per Pack | Order | | | |
|-----------------------|----------|----------------------------------|-------------|--|--|--|
| 30 tablet | \$ 2.33 | \$ 69.99 | Add to cart | | | |
| 60 tablet | \$ 2.17 | \$ 129.99 \$139.98 | Add to cart | | | |
| 90 tablet | \$ 2.11 | \$ 189.99 \$209.97 | Add to cart | | | |
| Free AirMail shipping | | | | | | |
| 120 tablet | \$ 1.92 | \$ 229.99 \$279.96 | Add to cart | | | |
| Free AirMail shipping | | | | | | |
| 180 tablet | \$ 1.78 | \$ 319.99 \$419.94 | Add to cart | | | |
| Free AirMail shipping | 9 | | | | | |

Gemfibrozil capsule

What is this medicine?

GEMFIBROZIL can help lower blood fats and cholesterol for people who are at risk of getting inflammation of the pancreas (pancreatitis) from having very high amounts of fats in their blood. This medicine is only for patients whose blood fats are not controlled by diet.

What should my health care professional know before I take this medicine?

They need to know if you have any of these conditions:

- cataracts
- · gallbladder disease
- heart disease
- · kidney or liver disease
- an unusual or allergic reaction to gemfibrozil, other medicines, foods, dyes, or preservatives
- · pregnant or trying to get pregnant
- · breast-feeding

How should I take this medicine?

Take this medicine by mouth with a glass of water. Take this medicine 30 minutes before a meal. Take your doses at regular intervals. Do not take your medicine more often than directed. Do not stop taking this medicine except on the advice of your doctor or health care professional.

Talk to your pediatrician regarding the use of this medicine in children. Special care may be needed.