

Desonate



Desonate is a steroid. It prevents the release of substances in the body that cause inflammation. **Desonate** topical (for the skin) is used to treat the inflammation and itching caused by a number of skin conditions such as allergic reactions, eczema, and psoriasis.

Active ingredient: [Desonide](#)

Brand(s): Dosetil Cream

Manufacturer: Glenmark Pharmaceuticals Ltd.

Disease(s): Inflammation / Itch / Allergy / Eczema / Psoriasis

Desonate 0.05% 10g

	Per pill	Per Pack	Order
1 tube	\$ 14.95	\$ 14.95	Add to cart
2 tube	\$ 12.47	\$ 24.95 \$29.90	Add to cart
3 tube	\$ 10.98	\$ 32.95 \$44.85	Add to cart
4 tube	\$ 9.99	\$ 39.95 \$59.80	Add to cart
5 tube	\$ 8.99	\$ 44.95 \$74.75	Add to cart

Desonide topical

What is this medicine?

DESONIDE is a steroid. It prevents the release of substances in the body that cause inflammation. Desonide topical (for the skin) is used to treat the inflammation and itching caused by a number of skin conditions such as allergic reactions, eczema, and psoriasis. Desonide topical may also be used for purposes not listed in this medication guide.

What should I tell my health care providers before I take this medicine?

You should not use desonide topical if you are allergic to it.

To make sure desonide topical is safe for you, tell your doctor if you have:

- any type of skin infection.

Also tell your doctor if you have diabetes. Steroid medicines may increase the glucose (sugar) levels in your blood or urine. You may also need to adjust the dose of your diabetes medications.

FDA pregnancy category C. It is not known whether desonide topical will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant while using this medicine.

It is not known whether desonide topical passes into breast milk or if it could harm a nursing baby. Tell your doctor if you are breast-feeding a baby.

Desonide topical is not approved for use by anyone younger than 18 years old. Children can absorb larger amounts of this medication through the skin and may be more likely to have unwanted side effects.

How should I use this medicine?

This medicine is usually applied 2 or 3 times daily. Follow all directions on your prescription label. Do not use this medicine in larger or smaller

