

## Compazine



Generic **Compazine** is used for controlling severe nausea and vomiting and treating schizophrenia. It is also used for the short-term (4 weeks) treatment of generalized nonpsychotic anxiety.

Active ingredient: [Prochlorperazine](#)

**Brand(s):**Vometil

**Manufacturer:** Cipla Limited

**Disease(s):**Nausea / Vomiting / Schizophrenia / Anxiety

### Compazine 5mg

	Per pill	Per Pack	Order
30 tablet	\$ 1.30	<b>\$ 38.95</b>	<a href="#">Add to cart</a>
60 tablet	\$ 1.18	<b>\$ 70.95</b> <del>\$77.90</del>	<a href="#">Add to cart</a>
90 tablet	\$ 1.09	<b>\$ 97.95</b> <del>\$116.85</del>	<a href="#">Add to cart</a>
120 tablet	\$ 0.98	<b>\$ 117.95</b> <del>\$155.80</del>	<a href="#">Add to cart</a>
180 tablet	\$ 0.88	<b>\$ 158.95</b> <del>\$233.70</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
360 tablet	\$ 0.78	<b>\$ 281.95</b> <del>\$467.40</del>	<a href="#">Add to cart</a>

Free AirMail shipping

## Prochlorperazine tablet

### What is this medicine?

PROCHLORPERAZINE helps to control severe nausea and vomiting. This medicine is also used to treat schizophrenia. It can also help patients who experience anxiety that is not due to psychological illness.

### What should I tell my health care provider before I take this medicine?

They need to know if you have any of these conditions:

- blood disorders or disease
- dementia
- liver disease or jaundice
- Parkinson's disease
- uncontrollable movement disorder
- an unusual or allergic reaction to prochlorperazine, other medicines, foods, dyes, or preservatives
- pregnant or trying to get pregnant
- breast-feeding

### How should I use this medicine?

Take this medicine by mouth with a glass of water. Take your doses at regular intervals. Do not take your medicine more often than directed. Do not stop taking this medicine suddenly. This can cause nausea, vomiting, and dizziness. Ask your doctor or health care professional for advice.

Talk to your pediatrician regarding the use of this medicine in children. Special care may be needed. While this drug may be prescribed for children as young as 2 years for selected conditions, precautions do apply.

