

Please click price below and you will be redirected to our pharmacy.

## Retrovir



Generic **Retrovir** is used for treating HIV infection when used along with other medicines. It is also used with other medicines to help prevent women from passing the HIV virus to the fetus during pregnancy.

Active ingredient: [Zidovudine](#)

**Brand(s):** Zidovir

**Manufacturer:** Cipla Limited

**Disease(s):** Hiv

### Retrovir 300mg

	Per pill	Per Pack	Order
10 tablet	\$ 7.00	<b>\$ 70.00</b>	<a href="#">Add to cart</a>
20 tablet	\$ 6.75	<b>\$ 135.00</b> <del>\$140.00</del>	<a href="#">Add to cart</a>
30 tablet	\$ 6.67	<b>\$ 200.00</b> <del>\$210.00</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
40 tablet	\$ 6.25	<b>\$ 250.00</b> <del>\$280.00</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
50 tablet	\$ 6.00	<b>\$ 300.00</b> <del>\$350.00</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
60 tablet	\$ 5.83	<b>\$ 350.00</b> <del>\$420.00</del>	<a href="#">Add to cart</a>
Free AirMail shipping			

## Zidovudine capsule

### What is this medicine?

ZIDOVUDINE ( ZDV) is an antiretroviral drug. It is used with other medicines to treat HIV. This medicine is not a cure for HIV. It will not stop the spread of HIV to others.

### What should I tell my health care provider before I take this medicine?

They need to know if you have any of these conditions:

- kidney disease
- liver disease
- low blood counts, like low white cell, platelet, or red cell counts
- an unusual or allergic reaction to zidovudine, other medicines, foods, dyes, or preservatives
- pregnant or trying to get pregnant
- breast-feeding

### How should I use this medicine?

Take this medicine by mouth with a full glass of water. You may take this medicine with or without food. Take your medicine at regular intervals. Do not take your medicine more often than directed. For your anti-HIV therapy to work as well as possible, take each dose exactly as prescribed. Do not skip doses or stop your medicine even if you feel better. Skipping doses may make the HIV virus resistant to this medicine and other

