#### **Esbriet**



Esbriet (pirfenidone) is used to treat a lung disease called idiopathic pulmonary fibrosis (IPF). IPF causes scar tissue to form deep within your lungs. The scar tissue thickens and becomes stiff or thick over time, which can make it harder for your lungs to work. Decreased lung function can make it hard for you to breathe.

Active ingredient: Pirfenidone

Brand(s):Pirfenex / Pifeheal

Manufacturer: Cipla LTd. / Healin Pharma Disease(s):Idiopathic Pulmonary Fibrosis

#### Esbriet 200mg

|                       | Per pill | Per Pack                         | Order       |  |  |  |
|-----------------------|----------|----------------------------------|-------------|--|--|--|
| 30 tablet             | \$ 2.00  | \$ 59.95                         | Add to cart |  |  |  |
| 60 tablet             | \$ 1.85  | \$ 110.95<br><del>\$119.90</del> | Add to cart |  |  |  |
| 90 tablet             | \$ 1.70  | \$ 152.95<br><del>\$179.85</del> | Add to cart |  |  |  |
| Free AirMail shipping |          |                                  |             |  |  |  |
| 120 tablet            | \$ 1.60  | \$ 191.95<br><del>\$239.80</del> | Add to cart |  |  |  |
| Free AirMail shipping |          |                                  |             |  |  |  |
| 180 tablet            | \$ 1.50  | \$ 269.95<br>\$359.70            | Add to cart |  |  |  |
| Free AirMail shipping | )        |                                  |             |  |  |  |

### Pirfenidone tablet

## What is pirfenidone?

Pirfenidone is used to treat a lung disease called idiopathic pulmonary fibrosis (IPF). IPF causes scar tissue to form deep within your lungs. The scar tissue thickens and becomes stiff or thick over time, which can make it harder for your lungs to work. Decreased lung function can make it hard for you to breathe. Other medical problems can occur when your brain, heart, and other organs do not get enough oxygen.

The cause of IPF is often unknown, but this condition is a progressive disease that can be fatal. Pirfenidone is not a cure for IPF, but pirfenidone may slow the progress of this disease.

Pirfenidone may also be used for purposes not listed in this medication quide.

# Important Information

Follow all directions on your medicine label and package. Tell each of your healthcare providers about all your medical conditions, allergies, and all medicines you use.

Before taking this medicine

You should not use pirfenidone if you are allergic to it.

To make sure pirfenidone is safe for you, tell your doctor if you have:

- · liver disease;
- · kidney disease; or
- · if you smoke.
- It is not known whether pirfenidone will harm an unborn baby. Tell
  your doctor if you are pregnant or plan to become pregnant.
- It is not known whether pirfenidone passes into breast milk or if it could affect the nursing baby. Tell your doctor if you are breast-