

Please click price below and you will be redirected to our pharmacy.

Revatio



Generic Revatio relaxes muscles and increases blood flow to particular areas of the body.

Active ingredient: [Sildenafil](#)

Brand(s):Assurans

Manufacturer: Cipla Limited

Disease(s):Pulmonary Arterial Hypertension

Revatio 20mg

	Per pill	Per Pack	Order
10 tablet	\$ 2.50	\$ 24.95	Add to cart
20 tablet	\$ 2.25	\$ 44.95 \$49.90	Add to cart
30 tablet	\$ 2.00	\$ 59.95 \$74.85	Add to cart
60 tablet	\$ 1.92	\$ 114.95 \$149.70	Add to cart
90 tablet	\$ 1.89	\$ 169.95 \$224.55	Add to cart
Free AirMail shipping			
120 tablet	\$ 1.83	\$ 219.95 \$299.40	Add to cart
Free AirMail shipping			
180 tablet	\$ 1.67	\$ 299.95 \$449.10	Add to cart
Free AirMail shipping			

Sildenafil

What is this medicine?

SILDENAFIL relaxes muscles and increases blood flow to particular areas of the body.

Sildenafil is used to treat pulmonary arterial hypertension and improve exercise capacity in men and women. Sildenafil under the brand name Viagra is used to treat erectile dysfunction (impotence) in men. Do not take Viagra while also taking Sildenafil, unless your doctor tells you to.

Sildenafil may also be used for other purposes not listed in this medication guide.

How should I use this medicine?

Do not take Sildenafil if you are also using a nitrate drug for chest pain or heart problems. This includes nitroglycerin (Nitrostat, Nitrolingual, Nitro-Dur, Nitro-Bid, and others), isosorbide dinitrate (Dilatrate-SR, Isordil, Sorbitrate), and isosorbide mononitrate (Imdur, ISMO, Monoket). Nitrates are also found in some recreational drugs such as amyl nitrate or nitrite ("poppers"). Taking Sildenafil with a nitrate medicine can cause a sudden and serious decrease in blood pressure. During sexual activity, if you become dizzy or nauseated, or have pain, numbness, or tingling in your chest, arms, neck, or jaw, stop and call your doctor right away. You could be having a serious side effect of Sildenafil. Do not take Sildenafil more than once a day. Allow 24 hours to pass between doses. Do not take Viagra while also taking Sildenafil, unless your doctor tells you to. Contact your doctor or seek emergency medical attention if your erection is painful or lasts longer than 4 hours. A prolonged erection (priapism) can

