

Buscopan



BUTYLSCOLOPAMINE (other names of active ingredient - scopolamine butylbromide, butylhyoscine and hyoscine butylbromide) is an anticholinergic medicine. Butylscopolamine has many effects in the body including decreasing the secretion of fluids, slowing the stomach and intestines, and dilation of the pupils. Butylscopolamine is used to relieve nausea, vomiting, and dizziness associated with motion sickness and recovery from anesthesia and surgery. Butylscopolamine may also be used in the treatment of parkinsonism, spastic muscle states, irritable bowel syndrome, diverticulitis, and other conditions.

Active ingredient: [Butylscopolamine](#)

Brand(s): Buscopan

Manufacturer: German Remedies Private Ltd.

Disease(s): Nausea / Vomiting / Dizziness / Sickness / Recovery From Anesthesia And Surgery / Parkinsonism / Irritable Bowel Syndrome / Diverticulitis / Spastic Muscle States

Buscopan 10mg

	Per pill	Per Pack	Order
10 tablet	\$ 0.99	\$ 9.95	Add to cart
20 tablet	\$ 0.85	\$ 16.95 \$19.90	Add to cart
30 tablet	\$ 0.83	\$ 24.95 \$29.85	Add to cart
60 tablet	\$ 0.80	\$ 47.95 \$59.70	Add to cart
90 tablet	\$ 0.78	\$ 69.95 \$89.55	Add to cart
180 tablet	\$ 0.75	\$ 134.95 \$179.10	Add to cart

Butylscopolamine tablet

What is this medicine?

BUTYLSCOLOPAMINE (other names of active ingredient - scopolamine butylbromide, butylhyoscine and hyoscine butylbromide) is an anticholinergic medicine. Butylscopolamine has many effects in the body including decreasing the secretion of fluids, slowing the stomach and intestines, and dilation of the pupils.

Butylscopolamine is used to relieve nausea, vomiting, and dizziness associated with motion sickness and recovery from anesthesia and surgery. Butylscopolamine may also be used in the treatment of parkinsonism, spastic muscle states, irritable bowel syndrome, diverticulitis, and other conditions.

Butylscopolamine may also be used for purposes other than those listed in this medication guide.

What should my health care professional know before I take this medicine?

Do not use Butylscopolamine without first talking to your doctor if you have

- kidney disease,
- liver disease,
- an enlarged prostate,
- difficulty urinating,

